



Vargas-Bello-Pérez, E., Bastías-Ruz, J., Toro-Mujica, P., Teixeira, D. L., & Enriquez-Hidalgo, D. (2020). Interplay between productive traits, the social rank, and the cow's stability in the order of entrance to the milking parlour. *Journal of Agricultural Science, Cambridge*, 158(6), 518-526. <https://doi.org/10.1017/S002185962000088X>

Peer reviewed version

License (if available):  
CC BY-NC-ND

Link to published version (if available):  
[10.1017/S002185962000088X](https://doi.org/10.1017/S002185962000088X)

[Link to publication record in Explore Bristol Research](#)  
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Cambridge University Press at <https://doi.org/10.1017/S002185962000088X>. Please refer to any applicable terms of use of the publisher.

## University of Bristol - Explore Bristol Research

### General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:  
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

# Interplay between productive traits, the social rank, and the cow's stability in the order of entrance to the milking parlour

E. Vargas-Bello-Pérez<sup>1,2,\*</sup>, J. Bastías-Ruz<sup>2</sup>, P. Toro-Mujica<sup>2,3</sup>, D.L. Teixeira<sup>2,3</sup>, D. Enriquez-Hidalgo<sup>2,4,5\*</sup>

<sup>1</sup>Department of Veterinary and Animal Sciences, Faculty of Health and Medical Sciences, University of Copenhagen, Grønnegårdsvej 3, DK-1870 Frederiksberg C, Denmark.

<sup>2</sup>Departamento de Ciencias Animales, Facultad de Agronomía e Ingeniería Forestal, Pontificia Universidad Católica de Chile, Santiago, Chile. Casilla-306. P. O. Box 6904411.

<sup>3</sup>Instituto de Ciencias Agronómicas y Veterinarias, Universidad de O'Higgins, Ruta 90 km 3, San Fernando, Chile.

<sup>4</sup>University of Bristol, Bristol Veterinary School, Langford, Somerset, BS40 5DU, UK

<sup>5</sup>Sustainable Agriculture Sciences Department, Rothamsted Research, North Wyke, Okehampton, Devon, EX20 2SB, United Kingdom

## \*Corresponding authors:

E. Vargas-Bello-Pérez [evargasb@sund.ku.dk](mailto:evargasb@sund.ku.dk) and D. Enriquez-Hidalgo [daniel.enriquez@uc.cl](mailto:daniel.enriquez@uc.cl)

## Acknowledgments

We would like to thank Fundación Agro UC for the animal facilities and assistance in obtaining the research data. Video recording equipment was kindly provided by Rafael Larraín and Jorge Oses. Special thanks should be given to Dr. Hanne Helen Hansen from the University of Copenhagen for proofreading our manuscript.

## Financial Support

This research received no specific grant from any funding agency, commercial or not-for-profit sectors.

## Conflicts of Interest

The authors declare there are no conflicts of interest.

28    **Ethical Standards**

29    Animal care and procedures were carried out according to the guidelines of the Animal Care and  
30    Use Committee of the Pontificia Universidad Católica de Chile (ID No. 150908002).

31

## **Interplay between productive traits, the social rank, and the cow's stability in the order of entrance to the milking parlour**

### **Abstract**

The aim of this study was to investigate whether social rank and stability in the order of entrance to the milking parlour are associated with production traits. The study was conducted on a dairy farm where cows ( $n = 215$ ) were managed in three groups according to lactation stage (Group 1: 78 cows 0-100 DIM; Group 2: 65 cows 101-200 DIM; and Group 3: 72 cows >200 DIM). Social rank (SR) was calculated from observations made from agonistic behaviour performed at the water troughs and feed bunks of each pen ( $n = 3$ ). The animals were classified in 3 levels of dominance based on at least 5 clear interactions, resulting in: 61 dominant, 75 intermediate and 69 subordinate cows based on SR. Stability in the order of entry was estimated as the standard deviation of the entry position. SR was weakly correlated to milk yield, urea and protein content in milk. The results showed that stable cows had higher milk production and entered the milking parlour after the non-stable animals. Stability in the order of entry to the milking parlour was not affected by social rank. Overall, the use of milking facilities appears to be associated with production traits rather than social rank.

**Keywords:** Social hierarchy, animal welfare, dairy, milking order.

## Introduction

Dairy cows form a social hierarchy through dominance establishment and this is associated with higher-ranking individuals having supremacy in the use of resources (Phillips and Rind, 2002). In a dairy herd, the social rank is related to factors such as age and animal size that can be measured as body condition score and live weight (Sottysiak and Nogalski, 2010). These factors are associated with physical strength and the competition for space that becomes a driver for aggressiveness of cows in confined spaces (Galindo and Broom, 2000). Aggressive social interactions create stress conditions between dominant and subordinate cows, which may affect milk production, especially for the subordinate cows (Chebel et al., 2016)

Dairy cows enter the milking area in a specific order and their behaviour is often expressed in a relatively consistent pattern (Paranhos da Costa & Broom 2001; Polikarpus et al., 2015). Consequently, preference for one side of the milking parlour seems to be a stable characteristic of a dairy cow (Hopster & van der Werf 1998; Grasso et al., 2007). Social rank (Melin et al. 2006), health-related characteristics (Flower et al. 2006; Polikarpus et al. 2015) and productive-related characteristics (Górecki & Wójtowski 2004) can influence the efficient use of milking facilities. For example, social rank, social structure of the herd and individual characteristics, such as anxiety, fear and stress, can influence side preference (Prelle et al. 2004). In addition, cows with locomotion and udder health problems can have problems to reach milking facilities (Gleeson et al., 2007).

The knowledge about milking order in dairy systems may provide data that can be used for early detection of health problems and prevention of production and economic losses (Polikarpus et al., 2015). Many studies have pointed out different factors affecting the use of milking facilities but to our knowledge, the interplay between productive traits, the social rank, and the stability of cows in the order of entrance to the milking parlour remain unexplored. Thus, the aim of this study was to determine whether social rank and stability in the order of entrance to the waiting area and milking parlour were affected by productive traits. To address the study's objective animals were initially classified by their social rank. We investigated the relationship between social rank and: (a) health related traits (locomotion score, body condition score and udder hygiene score); (b)

stability in the order of entrance to the waiting area and milking parlour; and (c) and milk production.

## **Materials and Methods**

### *Animals, housing and management*

Animal care and procedures were carried out according to the guidelines of the Animal Care and Use Committee of the Pontificia Universidad Católica de Chile (project ID No. 150908002). The study was conducted in a commercial dairy farm located in Pirque, Chile (33°38'28"S, 70°34'27"W). Lactating cows (n = 205; Holstein × Montbeliarde) were managed in three groups according to lactation stage (Group 1: 63 cows 0-100 DIM; Group 2: 77 cows 100-201 DIM; and Group 3: 65 cows >200 DIM). Animals were housed in dry lots (n=3; 80 × 80 m) with shade and had continuous access to water. Each pen had a rectangular-shape concrete water trough (3.5 × 1.5 m) and feed bunks were located in one side of the pens (approximately 60 linear meters per pen). Cows were fed three times per day (08:00, 10:00 and 17:00 h). Animals were milked in a 2 × 6 parallel milking parlour equipped with DELPRO™ farm manager system (DeLaval, Sweden), three times per day (03:30, 12:00 and 18:00 h). The walking distance from the pens to the milking parlour varied from 50 to 100m, depending on the group's location. Milk yield was recorded daily and analysed as total means. Body weights were individually measured after the first milking using a livestock chute with a weighing bar system. Body weights were recorded every 6 days and means were analysed. Milk components were recorded every 21 days (milk components were analysed by using an infrared analyser: Milko-Scan CombiFoss 6000; Foss Electric, Hillerød, Denmark) and means were analyzed.

### *Behavioural observations and social rank determination*

During 42 days, agonistic interactions (agonistic encounters included bunting, pushing, threatening, avoiding and fighting) were registered to calculate the social rank index of each animal and its subsequent correlation analysis with productive traits (milk production, somatic cell count, locomotion score, body condition score and udder hygiene score). In addition, the sequence of entrance to the waiting area and the milking parlour was used to define two indices of individual stability.

Behavioural observations were undertaken using a video recording system (SONY, YC-231G). Numbered coloured collars were used to facilitate animal identification. Agonistic interactions were registered through direct observation at the water trough after two milking times (12:00 and 18:00 h) during two consecutive days. Water trough access was denied from the moment the group left the pen to be milked until 30 minutes after the last cow entered to the pen (approximately one hour and 40 minutes without having access to water troughs in total). Each group was observed during five consecutive days for 30 hours. At the feed bunks, during two consecutive days, agonistic interactions were registered through direct observation two times per day (08:00 and 17:00 h). Each group was observed for 1 hour and 30 minutes by the same observer.

A social rank (SR) was calculated for each cow within each group from all observations (from water troughs and feed bunks). This SR was calculated based on methodology described by Galindo et al. (2000) and Hohenbrink and Meinecke-Tillmann (2012):  $\text{Social rank} = \text{interactions won} / (\text{won} + \text{lost interactions})$ . Animals showing a score of 0.0 to 0.4 were classified as subordinates. Individuals with a score from >0.4 to 0.6 were classified as intermediate. Cows with a score from >0.6 to 1.0 were classified as dominant.

#### *Health and productive related characteristics*

The locomotion score (LS), body condition score (BCS) and udder hygiene score (UHS) were considered as health and productive related characteristics. After behavioural observations, these characteristics were recorded at the end of the midday milking (12:00 h) during six consecutive days. The LS considered five individual aspects (spine curvature, tracking, speed, head carriage and abduction/adduction) of locomotion in a five-point scale as reported by O'Callaghan et al. (2003) and observations were done for once daily for 6 consecutive days. The BCS was based on a five-point scale where 1 = emaciated to 5 = overly fat (Wildman et al. 1982). A four-point score was used to measure the UHS based on Ruegg (2006) where 1= completely free of dirt or has very little dirt; 2= slightly dirty; 3 = mostly covered in dirt; and 4= completely covered, caked on dirt. The same person who was trained for those measurements carried out scorings for locomotion, body condition and udder hygiene.

*Milk production and stability in the order of entrance to the waiting area and milking parlour*

An automated management information system (DELPRO™, DeLaval, Sweden) recorded the entrance time to the milking parlour, milking duration, milking unit and individual milk yields and associates them with the ear tag embedded identification from each cow. Before each milking, all cows from each group were taken together to a waiting area. During this period, cows were free to choose their position in relation to the entrance in the milking parlour without any intervention of the stockman. The waiting area (120 m<sup>2</sup>) was in front of the milking parlour (Figure 1). Both sides of the milking parlour were identical and accessed through automatic gates. The gates were open as soon as a milking unit was empty.

The entrance order to the waiting area and the milking parlour were video-recorded (SONY, YC-231G). Nine records of entrance positions to the waiting area were obtained from each animal. For the entrance to the milking area 15 records of positions per animal were obtained. With these data, the individual stability of each animal was calculated as the standard deviation of its position records. With this information, 30% of the most stable cows and 30% of the less stable cows were selected to form two stability groups for production trait analyses. These groups (stable and non-stable) were compared for production characteristics such as milk production, urea, protein, fat and somatic cells in milk.

The sequence of entrance to the milking parlour was determined using the data obtained by the DelPro™ program (DeLaval) routinely used in the dairy farm, where the starting time of milking and milking unit used by each cow was recorded. Based on the different numbers of animals from Groups 1, 2 and 3, the positions in the order of entrance to the waiting area and milking parlour were standardize from 1 to 9.

*Statistical analysis*

Correlations were analysed using Pearson (parametric data: social rank, milk composition and milk yield) or Spearman (non-parametric data: locomotion score, body condition score, udder hygiene



and dominance level) tests. Additionally, Chi-square tests were performed to assess whether there was a dependency between Locomotion score (LS), body condition score (BCS) and udder hygiene score with social rank.

Productive characteristics, social rank, and order of entrance to the waiting area and to the milking parlour from stable and non-stable cows were analyzed by using ANOVA. The groups of cows (stable and non-stable) and their lactation stage were considered as fixed effects in the following model:

$$Y_{ijk} = \mu + \alpha_i + \beta_k + (\alpha\beta)_{ij} + \varepsilon_{ijk}$$

Where  $Y_{ijk}$  is the dependent variable;  $\mu$  is the overall mean;  $\alpha_i$  is the effect of lactation stage;  $\beta_k$  is the effect of stability group; and  $\varepsilon_{ijk}$  is the experimental error.

The homogeneity of variances was analysed with the Levene test. This was followed by the Student-Newman-Keuls multiple comparison of means procedure at  $P < 0.05$  to determine differences between the different groups. Results were expressed as mean  $\pm$  standard deviation. Probability of  $P < 0.05$  was defined as significant. The SPSS statistical software for Windows was used (version 15.0.0; SPSS Inc., Chicago IL, USA). When statistical analyses were performed, the outliers identified were identified as the consequence of extraordinary events, such as health problems (i.e., clinical mastitis and abortion), and the need for some veterinary treatment. Therefore, as they did not represent an important segment of the study population, they were eliminated from the analyses.

Additionally, to assess whether there was an effect of position on the standard deviation of the position data, the cows were grouped into 9 groups according to the average position obtained during the observations, and the SD of their locations was calculated. ANOVA or Kruskal-Wallis test were performed with these data, after heteroscedasticity test of variances. When there was no heteroscedasticity of variance, the multiple comparison was performed using the Tukey's

Honest Significant Difference test, and otherwise, through the non-parametric test by Notched Box Plots.

## Results

Out of the initial 215 cows in the herd, five cows were culled out due to health problems during the study period; one additional cow had an abortion and was discarded from the study due to the need for veterinary treatment. Of the remaining 209 cows, 4 were eliminated from the results database, because they presented outlier values in some of the variables analyzed. Table 1 shows the final production and health related characteristics for each group. Briefly, group 1 consisted of 77 animals, of which 27% were dominant, 38% were intermediate and 35% were subordinate. Group 2 (n = 63) had a distribution of 32% dominant animals, 29% of intermediate animals and 38% of subordinate animals. Group 3 (n = 65), where the animals in their last third of lactation had a distribution of 30% of dominant animals, 39% of intermediate animals and 31% of subordinate animals. The social rank evaluation from all animals (n = 205) resulted in 61 dominant animals, 75 intermediate animals and 69 subordinate animals. The agonistic interactions during water consumption represented less than 20% of the total interactions (Table 2).

There was a significant correlation between the SR and parity, which means that multiparous cows tend to be dominant individuals in the herd (Table 3). A weak correlation between parity and locomotion score ( $r = 0.464$ ,  $P < 0.001$ ) was found, which in a way indicates that animals with more parities have higher locomotion scores. Body condition score had a tendency to be weakly correlated with the dominant animals, where dominant animals had better scores than the subordinate animals ( $r = 0.126$ ,  $P = 0.095$ ). Table 4 shows the distribution of locomotion score, body condition score and udder hygiene score of cows within social ranks. Regardless of social rank in the herd, 88% of the cows had a locomotion score of 2, 86% of the cows had a body condition score between 2.75 and 3.25 and 78% of the cows had a udder hygiene score between 1 and 2. The Chi-square test showed that there was no dependency between locomotion score [X<sup>2</sup> (205) 6, 3.94,  $P = 0.689$ ], body condition score [X<sup>2</sup> (205) 10, 14.3,  $P = 0.155$ ] and udder hygiene score [X<sup>2</sup> (205) 6, 4.3,  $P = 0.634$ ] on social rank (Table 4).

The stability in the order of entrance to the waiting area had a significant effect on milk production, thus, stable cows had higher milk production than non-stable cows (Table 5). In relation to milk fat, stable cows presented a higher content at the waiting area and in the milking parlour, however, significant differences were only observed in the milking parlour. The stability in the order of entrance also affected the concentration of urea at the waiting area with higher contents detected in stable cows.

As shown in Figure 2, the standard deviation of the position of the cows was related to their average position, thus, at the waiting room, the ANOVA test was significant, and differences were observed between the standard deviations of each category. Thus, the Kruskal-Wallis test was used, which gave a value of  $P = <0.001$ . This test showed differences between categories 1, 2 and 9, which presented the smallest standard deviations, with the central groups from 4 to 7. At the milking parlour, the ANOVA test was not significant ( $P = 0.068$ ) and, as in the waiting room, the standard deviation of the data presented heteroscedasticity, so the Kruskal-Wallis test was used, which was significant ( $P = 0.004$ ), obtaining differences between the means of groups 1, 2 and 8 with group 4.

## Discussion

### *Productive traits and social rank*

In this study, the social rank was determined on agonistic interactions from each group that was observed during the first minutes when there was no access to water. These findings coincide with a study conducted by Kondo et al. (1984) who observed a rapid decrease in the frequency of agonistic encounters as time passes. Similarly, at the feed bunk, Menke et al. (2000) reported that agonistic behaviour decreases in a relative short period.

Normally, the hierarchy is expressed mainly at the feed bunks (Val-Laillet et al. 2008). In this study, cows had water *ad libitum* after the 1:40 min restriction imposed after milking and feed was offered three times a day. In this regard, feed availability was restricted creating more competition among animals, resulting in a greater number of agonistic interactions at the time of feeding (Proudfoot et al. 2009). The results of a study conducted by DeVries and Von Keyserlingk (2006),

concluded that dairy cows experience a lower number of agonistic interactions when they are given more than the standard of 0.61 m of feeder/animal (Grant & Albright 2001). In the current study, animals had 0.8 m/animal. Previous studies showed that fresh cows, first calving heifers and cows recently integrated into a group are submissive cows; and larger and older cows in a group are often the most dominant (Guhl & Atkeson 1959; Grant & Albright 2001; Phillips & Rind 2002).

The correlations between locomotion scores and parity found in this study were weak, however they pointed at the fact that older animals are more prone to having foot problems (Galindo & Broom 2000). Hetticha et al. (2007) reported that the number of births and, therefore, the age, was a risk factor for the presentation of this pathology, where cows with three or more lactations presented 3.8 times more chances of suffering from this condition. This agrees with Sogstad et al. (2005), who found that cows with three or more lactations had higher risks of lameness than those with less than three lactations. In this study, animals that are more dominant had increased body conditions scores. This finding suggest that dominant animals have unrestricted access to different resources (i.e., feed, water, and bedding), while subordinate animals have restricted access to the feeding places, which could affect their feed intake (Dickson et al. 1970; Llonch et al., 2018) and, therefore, their body condition score. Overall, in this study, given the weak correlations and non-significant Chi-square values found between locomotion score, body condition score, udder hygiene score and social rank. These results therefore, should be interpreted with caution as this was based on observations from 205 animals, which was not sufficient to have a greater dispersion of scores.

There was a weak correlation between milk yield (mean milk yield for each cow) and dominance; dominant animals had higher performance than the subordinate animals ( $r = 0.191$ ,  $P < 0.001$ ). In dry-lot pens, where feeding is not available *ad libitum*, animals unable to adapt to a competitive environment may be at a disadvantage in terms of the quantity and quality of the feed they have access to (Zobel et al. 2011), which could affect their consumption and, therefore, decrease their production. It is very important that the cows have good access to the diet and water throughout the day, to ensure the health, welfare, production and efficiency of the cow (Shabi et al. 2005). In farms where feed is limited, and competition is high, dominant cows eat more feed (Grant & Albright 2001) and drink more water (Teixeira et al. 2009) than subordinate cows.

291

292 There were weak correlations between social rank and body weight ( $r = 0.276$ ,  $P < 0.001$ ), parity ( $r$   
293  $= 0.339$ ,  $P < 0.001$ ) and milk yield ( $r = 0.191$ ,  $P < 0.001$ ). In this study, when social rank was  
294 determined at the water troughs and feed bunks, it was observed that dominant animals with  
295 larger body size, had unimpeded access to feed and water, while the subordinates were displaced  
296 and forced to move to other places without being able to consume feed. These results coincided  
297 with the study on a single herd carried out by Dickson et al. (1970), who suggested that live body  
298 weights and age were the factors most closely related to the position of the cow in the social  
299 hierarchy. In addition, Schein and Fohrman (1955) reported a high significant relationship between  
300 the range of dominance and age, and live body weight as well. On the other hand, the correlation  
301 between the social rank and body condition score and udder hygiene were low; the correlation  
302 coefficients were between  $0.096$  and  $0.126$  (Table 3). No significant correlation was observed  
303 between SR and locomotion scores nor somatic cell count (Table 3).

304

305 *Stability in the order of entrance to the waiting area and milking parlour*

306 *At the waiting area*, we observed a relationship ( $P = 0.008$ ) between SR and milk yield, since stable  
307 cows (lower standard deviation) were those with higher milk yield, compared to non-stable cows  
308 (higher standard deviation) (Table 5). Furthermore, although stable cows entered into the waiting  
309 area 0.12 places after non-stable cows (Table 5) this is not expected to be biologically significant.  
310 According to (Grasso et al. 2007) the cows have a coherent order of entry to the waiting area,  
311 which is considered as a social characteristic of the animals belonging to a dairy system.

312

313 The milk urea contents in both groups are within the normal expected range (150 to 420 mg/L)  
314 (Westwood et al. 1998). Stable cows had higher contents of milk urea. The concentration of milk  
315 urea is related to the supply of nitrogen in the diet, specifically the balance between degradable  
316 proteins and energy in the rumen. A high nitrogen intake or an energy deficiency produce an  
317 excess of ammonium in the rumen, which is absorbed and transformed into urea in the liver. This  
318 increases its concentration in both blood and milk, and its excretion in the urine. These  
319 concentrations are important since they can have repercussions for the health, fertility or  
320 efficiency in the milk production of the animal (Noro & Wittwer 2003).

We observed a relationship ( $P < 0.001$ ) between the stability in the order of entrance to the milking parlour and milk fat contents (Table 5). There were no studies found on the association that might exist between the stability in the order of entrance to milking facilities and milk composition. It is known that breed, parity and stage of lactation affect milk yield and milk composition (Kelsey et al. 2003). In this study, we were not able to select animals based on parity or number of calvings, which could be helpful to explain the observed relationship.

In relation to the stability in the order of entrance to the milking parlour, the stable animals entered 1.7 positions after non stable animals, which agree with Hopster and van der Werf (1998) who found that cows with a consistent side choice took more time to enter the room. Rathore (1982) found that high-yield cows voluntarily entered the milking parlour earlier than low-yielding cows, and that was explained by the fact that cows were relieving the pressure of the udder caused by the milk, and that was the source of motivation to enter the milking parlour early. However, in the current study, we did not observed a relationship between the stability in the order of entrance to the milking parlour and milk yield. Conversely, Grasso et al. (2007) observed a positive correlation between milking order and milk production in primiparous cows, although the correlation coefficient was not high ( $r = 0.22$ ). As shown in Table 5, there were no interactions between lactation stage (G) and stability (S) for milk yield, which could indicate that stability is a characteristic associated with the animal independently from its days in milk.

Overall, results showed that a cow that prefers to come in first place would likely have less variation in her milking position than a cow in the middle of the herd because it may be easier to consistently be first than consistently be last. Also, because there is less scope for variation at each end of the milking order. Although, our study used 205 animals, the heteroscedastic nature of milking position variance in the milking order was also reported in larger herds of around 500 lactating cows (Beggs et al., 2018).

*Limitations of the study*

Additional factors need to be taken into account when interpreting the data from this study. The weak correlations obtained are from the total available animals, without differentiating them by production level. Thus, there was a high percentage of the variance that is a consequence of other factors, some of which were considered in the analysis (lactation group and body weight) and others that the study did not account, such as the number of calvings and months of gestation. In this sense, the significance of the correlations could indicate that, by decreasing the sources of variation in the analyzed variables, the value of their correlation with social dominance could increase. Thus, although correlation values weak, they could be considered as the basis for a larger controlled experimental design in which the effect of the aforementioned variables are included or isolated.

## Conclusion

In the current study, social rank was weakly correlated to production parameters such as milk yield, protein content, urea content, parity and body weight. Social rank did not affect cow's stability in the order of entrance to the waiting area and milking parlour. Stable cows had higher milk production and entered the milking parlour after non-stable animals. Overall, under the conditions of this study, the stability of the milking order appeared to be more closely associated with production traits rather than social rank.

## References

- BEGGS, D. S., E. C. JONGMAN, P. H. HEMSWORTH, AND A. D. FISHER. 2018. SHORT COMMUNICATION: MILKING ORDER CONSISTENCY OF DAIRY COWS IN LARGE AUSTRALIAN HERDS. *JOURNAL OF DAIRY SCIENCE* 101(1):603-608.
- CHEBEL, R. C., SILVA, P. R. B., ENDRES, M. I., BALLOU, M. A., LUCHTERHAND, K. L., (2016). SOCIAL STRESSORS AND THEIR EFFECTS ON IMMUNITY AND HEALTH OF PERIPARTURIENT DAIRY COWS. *J. DAIRY SCI.* 99, 3217–3228.
- DEVRIES, T. & VON KEYSERLINGK, M. (2006). Feed stalls affect the social and feeding behavior of lactating dairy cows. *Journal of Dairy Science* 89(9), 3522-3531.
- DICKSON, D. BARR, G. JOHNSON, L. & WIECKERT, D. (1970). Social dominance and temperament of holstein Cows. *Journal of Dairy Science* 53(7), 904-907.
- FLOWER, F. SANDERSON, D. & WEARY, D. (2006). Effects of milking on dairy cow gait. *Journal of Dairy Science* 89(6), 2084-2089.
- GALINDO, F. & BROOM, D. (2000). The relationships between social behaviour of dairy cows and the occurrence of lameness in three herds. *Veterinary Science* 69(1), 75-79.
- GLEESON, D. E. O'BRIEN, B. BOYLE, L. & EARLY, B. (2007). Effect of milking frequency and nutritional level on aspects of the health and welfare of dairy cows. *Animal* 1, 125-132.
- GÓRECKI, M. T. & WÓJTOWSKI, J. (2004). Stability of milking order in goat over a long period. *Archives Animal Breeding* 47(2), 203-208.

- GRANT, R. J. & ALBRIGHT, J. L. (2001). Effect of animal grouping on feeding behavior and intake of dairy cattle. *Journal of Dairy Science* **84**, 156-163.
- GRASSO, F. DE ROSA, G. NAPOLITANO, F. DI FRANCIA, A. & BORDI, A. (2007). Entrance order and side preference of dairy cows in the milking parlour. *Italian Journal of Animal Science* **6**(2), 187-194.
- GUHL, A. & ATKESON, F. (1959). Social organization in a herd of dairy cows. *Kansas Academy of Science* **62**(1), 80-87.
- HETTICHA, E. HINOSTROZAB, M. VAN SCHAIKB, G. & TADICH, N. (2007). Factors associated to lameness in 50 dairy herds in the Xth Region, Chile. *Archivos médicos veterinarios* **39**(3), 247-253.
- HOHENBRINK, S. & MEINECKE-TILLMANN, S. (2012). Influence of social dominance on the secondary sex ratio and factors affecting hierarchy in Holstein dairy cows. *Journal of Dairy Science* **95**(10), 5694-5701.
- HOPSTER, H. & VAN DER WERF, J. T. (1998). Side preference of dairy cows in the milking parlour and its effects on behaviour and heart rate during milking. *Applied Animal Behaviour Science* **55**(3-4), 213-229.
- KELSEY, J. A. CORL, B. A. COLLIER, R. J. & BAUMAN, D. E. (2003). The effect of breed, parity, and stage of lactation on conjugated linoleic acid (CLA) in milk fat from dairy cows. *Journal of Dairy Science* **86**(8), 2588-2597.
- LLONCH, P., MAINAU, E., IPHARRAGUERRE, I. R., BARGO, F., TEDÓ, G., BLANCH, M., & MANTECA, X. (2018). Chicken or the Egg: The Reciprocal Association Between Feeding Behavior and Animal Welfare and Their Impact on Productivity in Dairy Cows. *Frontiers in veterinary science*, 5, 305.
- MENKE C, WAIBLINGER S, FÖLSCH DW. (2000) Die Bedeutung von Managementmassnahmen im Laufstall für das Sozialverhalten von Milchkühen [Importance of herd management in loose housing systems in the social behavior of dairy cows]. *DTSCH TIERARZTL WOCHENSCHR.*; **107**(7), 262-268.
- MELIN, M. HERMANS, G. PETTERSSON, G. & WIKTORSSON, H. (2006). Cow traffic in relation to social rank and motivation of cows in an automatic milking system with control gates and an open waiting area. *Applied Animal Behaviour Science* **96**(3-4), 201-214.
- NORO, M. & WITTWER, F. (2003). Utilidad de la determinación de la urea en la leche. *Vetermás* **2**, 2-5.
- O'CALLAGHAN, K. A., P. J. CRIPPS, D. Y. DOWNHAM, AND R. D. MURRAY. (2003). SUBJECTIVE AND OBJECTIVE ASSESSMENT OF PAIN AND DISCOMFORT DUE TO LAMENESS IN DAIRY CATTLE. *ANIMAL WELFARE* **12**,(4), 605-610.
- PARANHOS DA COSTA, M. & BROOM, D. (2001). Consistency of side choice in the milking parlour by Holstein–Friesian cows and its relationship with their reactivity and milk yield. *Applied Animal Behaviour Science* **70**(3), 177-186.
- PHILLIPS, C. & RIND, M. (2002). The effects of social dominance on the production and behavior of grazing dairy cows offered forage supplements. *Journal of Dairy Science* **85**(1), 51-59.
- POLIKARPUS, A. KAART, T. MOOTSE, H. DE ROSA, G. & ARNEY, D. (2015). Influences of various factors on cows' entrance order into the milking parlour. *Applied Animal Behaviour Science* **166**, 20-24.
- PRELLE, I. PHILLIPS, C. DA COSTA, M. P. VANDENBERGHE, N. & BROOM, D. (2004). Are cows that consistently enter the same side of a two-sided milking parlour more fearful of novel situations or more competitive? *Applied Animal Behaviour Science* **87**(3-4), 193-203.
- PROUDFOOT, K. L. VEIRA, D. WEARY, D. & VON KEYSERLINGK, M. (2009). Competition at the feed bunk changes the feeding, standing, and social behavior of transition dairy cows. *Journal of Dairy Science* **92**(7), 3116-3123.



- RATHORE, A. (1982). Order of cow entry at milking and its relationships with milk yield and consistency of the order. *Applied Animal Behaviour Science* **8**(1), 45-52.
- RUEGG, P. L. (2006). The role of hygiene in efficient milking. In *Western Canadian Dairy Seminar (WCDS): Adv. Dairy Tech.* pp. 285-293.
- SCHEIN, M. W. & FOHRMAN, M. H. (1955). Social dominance relationships in a herd of dairy cattle. *The British Journal of Animal Behaviour* **3**(2), 45-55.
- SHABI, Z. MURPHY, M. & MOALLEM, U. (2005). Within-day feeding behavior of lactating dairy cows measured using a real-time control system. *Journal of Dairy Science* **88**(5), 1848-1854.
- SOGSTAD, Å. FJELDAAS, T. & ØSTERÅS, O. (2005). Lameness and claw lesions of the Norwegian red dairy cattle housed in free stalls in relation to environment, parity and stage of lactation. *Acta Veterinaria Scandinavica* **46**(4), 203.
- SOTTYSIAK, T. NOGALSKI, Z. (2010). THE EFFECTS OF SOCIAL HIERARCHY IN A DAIRY CATTLE HERD ON MILK YIELD. *POLISH JOURNAL OF NATURAL SCIENCES* **25**: 22-23
- TEIXEIRA, D. L. HÖTZEL, M. J. MACHADO FILHO, L. C. P. ENRÍQUEZ HIDALGO, D. & CAZALE, J. D. (2009). Aspectos etológicos no suprimento de água em bovinos leiteiros. *Biotemas* **22**(4), 193-198.
- VAL-LAILLET, D. DE PASSILLÉ, A. M. RUSHEN, J. & VON KEYSERLINGK, M. A. (2008). The concept of social dominance and the social distribution of feeding-related displacements between cows. *Applied Animal Behaviour Science* **111**(1-2), 158-172.
- WESTWOOD, C. LEAN, I. & KELLAWAY, R. (1998). Indications and implications for testing of milk urea in dairy cattle: a quantitative review. Part 1. Dietary protein sources and metabolism. *New Zealand Veterinary Journal* **46**(3), 87-96.
- WILDMAN, E. JONES, G. WAGNER, P. BOMAN, R. TROUTT JR, H. & LESCH, T. (1982). A dairy cow body condition scoring system and its relationship to selected production characteristics. *Journal of Dairy Science* **65**(3), 495-501.
- ZOBEL, G. SCHWARTZKOPF-GENSWEIN, K. GENSWEIN, B. & VON KEYSERLINGK, M. (2011). Impact of agonistic interactions on feeding behaviours when beef heifers are fed in a competitive feeding environment. *Livestock Science* **137**(1-3), 1-9.

**Table 1.** Productive parameters, health related characteristics and social ranks (mean  $\pm$  standard error)

Parameter	Group 1 (n = 77)	Group 2 (n = 63)	Group 3 (n = 65)
	0 – 100 days in milk	101 – 200 days in milk	>200 days in milk
Productive characteristics			
Milk yield, kg/d	31.6 $\pm$ 0.69	34.6 $\pm$ 0.52	34.0 $\pm$ 0.62
Fat, g/kg	37.5 $\pm$ 0.70	36.8 $\pm$ 0.80	34.8 $\pm$ 0.60
Protein, g/kg	32.2 $\pm$ 0.20	32.7 $\pm$ 0.40	32.9 $\pm$ 0.30
Milk urea N, mg/L	346 $\pm$ 9.6	366 $\pm$ 6.1	387 $\pm$ 5.6
Somatic cell counts, $\times 10^3$ mL	210 $\pm$ 44.6	198 $\pm$ 30.1	207 $\pm$ 38.3
Live body weight, kg	622 $\pm$ 12.8	662 $\pm$ 12.0	700 $\pm$ 9.4
Number of lactations	2.06 $\pm$ 0.170	2.39 $\pm$ 0.170	1.86 $\pm$ 0.150
Health related characteristics			
Locomotion score	2.17 $\pm$ 0.050	2.08 $\pm$ 0.040	2.12 $\pm$ 0.040
Body condition score	3.07 $\pm$ 0.030	2.95 $\pm$ 0.020	3.10 $\pm$ 0.020
Udder hygiene score	1.65 $\pm$ 0.090	1.87 $\pm$ 0.080	2.03 $\pm$ 0.860

466 **Table 2.** Agonistic interactions recorded for 5 consecutive days in 205 dairy cows

Agonistic interactions	Group 1 (n = 77)	Group 2 (n = 63)	Group 3 (n = 65)
At the water troughs	232	178	147
At the feed bunks	800	630	685
Total interactions	1032	808	832
Social rank, number of cows			
Dominant	21	21	20
Intermediate	29	18	25
Subordinate	27	24	20

467

468

469 **Table 3.** Correlations between different parameters for animals with at least 5 encounters

Parameters	Correlation (R)	P-value
Parity and locomotion score	0.464	<0.001
Milk yield and body weight	0.445	<0.001
Milk yield and somatic cell count	- 0.101	0.308
Body condition score and locomotion score	- 0.016	0.824
Social rank and parity	0.339	<0.001
Social rank and milk protein content	0.293	<0.001
Social rank and body weight	0.276	<0.001
Social rank and milk urea content	0.242	<0.001
Social rank and milk yield	0.191	<0.001
Social rank and body condition score	0.126	0.095
Social rank and udder hygiene score	0.096	0.083
Social rank and locomotion score	0.056	0.518
Social rank and somatic cell count	-0.023	0.738
Social rank and milk fat content	- 0.126	0.056

470 Correlations were analysed using Pearson (parametric data: social rank, milk composition and milk  
471 yield) or Spearman (non-parametric data: locomotion score, body condition score, udder hygiene  
472 and dominance level) tests.

473  
474  
475  
476

477 **Table 4.** Distribution of locomotion score (LS), body condition score (BCS) and udder hygiene score  
478 (UHS) of lactating cows in relation to their social rank and Chi-square analysis

Social rank								
Items	Dominant		Intermediate		Subordinate		All herd	
	n	%	n	%	n	%	n	%
LS								
LS 1	0	0	1	1	0	0	1	0.5
LS 2	54	89	63	84	62	90	179	88
LS 3	7	11	10	13	7	10	24	11
LS 4	0	0	1	1	0	0	1	0.5
LS 5	0	0	0	0	0	0	0	0
Cows <sub>total</sub>	61		75		69		205	
BCS								
BCS 2.50	1	2	1	1	2	4	4	3
BCS 2.75	13	21	18	24	16	22	47	22
BCS 3.00	22	35	34	45	33	45	89	42
BCS 3.25	13	23	14	19	16	26	43	22
BCS 3.50	9	15	8	11	2	3	19	9
BCS 3.75	3	5	0	0	0	0	3	2
Cows <sub>total</sub>	61		75		69		205	
UHS								
UHS 1	17	29	32	43	26	37	75	37
UHS 2	28	45	27	36	28	41	83	41
UHS 3	15	24	16	21	14	19	45	21
UHS 4	1	2	0	0	1	3	2	1
Cows <sub>total</sub>	61		75		69		205	
LS	Dominant	Intermediate	Subordinate	X <sup>2</sup>				
LS 1	0	1	0			Chi-square	3.94	
LS 2	54	63	62			Degrees of freedom	6	
LS 3	7	10	7	0.68	P-value		0.689	
LS 4	0	1	0					
LS 5	0	0	0					
BCS	Dominant	Intermediate	Subordinate	X <sup>2</sup>				
BCS 2.50	1	1	2			Chi-square	14.3	
BCS 2.75	13	18	16			Degrees of freedom	10	
BCS 3.00	22	34	33	0.16	P-value		0.155	
BCS 3.25	13	14	16					
BCS 3.50	9	8	2					
BCS 3.75	3	0	0					
UHS	Dominant	Intermediate	Subordinate	X <sup>2</sup>				
UHS 1	17	32	26			Chi-square	4.3	
UHS 2	28	27	28			Degrees of freedom	6	
UHS 3	15	16	14	0.63	P-value		0.634	
UHS 4	1	0	1					

479  
480


---

**Table 5.** Productive characteristics, social rank, and average order of entrance to the milking parlour from stable and no stable cows (mean  $\pm$  standard deviation)

Parameters	Stable cows	No stable cows	<i>P</i> -value (G)	<i>P</i> -value (S)	<i>P</i> -value (G×S)
Waiting area					
Milk yield, kg/d	34.1 $\pm$ 0.67	32.3 $\pm$ 0.73	0.008	0.036	0.316
Fat, g/kg	36.0 $\pm$ 0.90	37.0 $\pm$ 0.70	0.155	0.362	0.818
Protein, g/kg	32.0 $\pm$ 0.30	32.0 $\pm$ 0.30	0.223	0.328	0.837
Milk urea N, mg/L	378 $\pm$ 6.7	351 $\pm$ 8.8	0.007	0.005	0.293
Somatic cell count, $\times 10^3$ /ml	169 $\pm$ 34.9	186 $\pm$ 33.4	0.291	0.994	0.396
Social rank	0.5 $\pm$ 0.02	0.5 $\pm$ 0.02	0.786	0.203	0.210
Order of entrance <sup>1</sup>	4.6 $\pm$ 0.28	4.7 $\pm$ 0.16	<0.001	0.637	0.799
Milking parlour					
Milk yield, kg/d	34.6 $\pm$ 0.61	33.7 $\pm$ 0.68	0.079	0.283	0.312
Fat, g/kg	36.5 $\pm$ 0.80	37.1 $\pm$ 0.70	0.489	<0.001	0.622
Protein, g/kg	32.4 $\pm$ 0.30	32.0 $\pm$ 0.30	0.153	0.491	0.860
Milk urea N, mg/L	371 $\pm$ 7.4	365 $\pm$ 8.5	0.005	0.618	0.569
Somatic cell count, $\times 10^3$ /ml	190 $\pm$ 37.9	202 $\pm$ 33.7	0.423	0.928	0.065
Social rank	0.5 $\pm$ 0.22	0.5 $\pm$ 0.24	0.703	0.619	0.644
Order of entrance <sup>1</sup>	6.9 $\pm$ 0.37	5.2 $\pm$ 0.18	<0.001	<0.001	<0.001

G = probability of lactation stage effect; S = probability of stability effect.

<sup>1</sup>Based on the different numbers of animals from Groups 1, 2 and 3, the positions in the order of entrance to the waiting area and milking parlour were standardize from 1 to 9.




**Figure 1.** Layout of the 2 × 6 parallel milking parlour. 1-entrance, 2-waiting area, 3-operator's pit, 4-right alley, 5-left alley, 6-exit alley.



506


507

508



509

510



**Figure 2.** Notched Box Plots from average order of entrance to the waiting area and milking parlour